

# Use of Microwave Assited Extraction to obtain anthocyanins and total phenolic compounds rich extracts in myrtle (*Myrtus communis* L.)

Ana V. González de Peredo<sup>1</sup>, Gerardo F. Barbero<sup>1,\*</sup>, Ana Jiménez-Cantizano<sup>2</sup>, Ceferino Carrera<sup>3</sup>, Miguel Palma<sup>1</sup> and Carmelo G. Barroso<sup>1</sup> \*(gerardo.fernandez@uca.es)

<sup>1</sup> Department of Analytical Chemistry, Faculty of Sciences, IVAGRO, ceiA3, University of Cadiz. Av. República Saharaui s/n. 11510, Puerto Real, Cadiz, Spain.


<sup>2</sup> Department of Chemical Engineering and Food Technology, Faculty of Sciences, IVAGRO, ceiA3, Av. República Saharaui s/n. 11510, Puerto Real, Cadiz, Spain.

<sup>3</sup> Instituto de Investigación Vitivinícola y Agroalimentario (IVAGRO), ceiA3, University of Cadiz, Av. República Saharaui s/n. 11510, Puerto Real, Cadiz, Spain.



## INTRODUCTION

- ❖ Myrtle is the fruit of the Myrtle shrub (*Myrtus communis* L.).<sup>1</sup>
- ❖ It belongs to the genus *Myrtus* and the family *Myrtaceae*.
- ❖ It grows spontaneously in the Mediterranean area and the Middle East.<sup>2</sup>
- ❖ This species is a very aromatic plant because of the high essential oil content in its leaf, flower and fruit glands.<sup>3</sup>
- ❖ It contains a huge concentration of antioxidant substances (anthocyanins and phenolic compounds), much higher than other fruits.<sup>4</sup>
- ❖ It is used to produce the characteristic myrtle liqueur typical of Sardinia.<sup>5</sup>
- ❖ Anthocyanins and phenolics have potential health-promoting effects (i.e., antioxidant, anti-inflammatory, and anticancer activities).<sup>6</sup>



## MATERIALS AND METHODS

### MICROWAVE ASSISTED EXTRACTION



❖ Extraction time: 5 min



- ❖ Microwave MARS 6 240/50 (one touch technology, CEM Corporation Matthews, North Carolina, United State).
- ❖ Power: 800 W.



❖ Keep at -20 °C until analysis


## RESULTS AND DISCUSSION

### EXPERIMENTAL DESIGN (BOX BEHNKEN)


27 duplicate extractions (1 block)  
4 Factors (3 Levels):

%MeOH	pH	Temperature (°C)	Ratio (mL)
25 – 50 – 75	2 – 4,5 – 7	10 - 35-60	10 – 15 – 20


2 Responses: Total anthocyanins and total phenolics



### TOTAL PHENOLICS




### Extraction time





### Optimum condition


% MeOH in water	pH	Temperature (°C)	Ratio	Time (5 min)
59.64	2	100	20	15



### ANTHOCYANINS



### Extraction time



### Optimum condition

% MeOH in water	pH	Temperature (°C)	Ratio	Time (5 min)
50	3,33	50	20	2

### CONCLUSIONS

- ❖ Two methods for the extraction of total anthocyanins and total phenolic compounds in myrtle by Microwave Assited Extraction have been developed.
- ❖ A Box-Behnken experimental design with 4 factors and 2 responses (one for anthocyanins and one for phenolic compounds) has been carried out.
- ❖ HPLC has been used for the identification and analysis of the anthocyanins, and Folin Cioacalteu method was used for the analysis of total phenolic compounds.
- ❖ According to the experimental design, the percentage of methanol in the extraction solvent, the interaction percentage of methanol in the extraction solvent-percentage of methanol in the extraction solvent and the pH are the determining factors for the extraction of total phenolic compounds.
- ❖ According to the experimental design, the interaction percentage of methanol in the extraction solvent-percentage of methanol in the extraction solvent, temperatura and the interaction pH-temperature are the determining factors for the extraction of anthocyanins.
- ❖ Different extraction times were tested using the optimal conditions of extraction. A extraction time of 15 minutes was enough for quantitative extraction of total phenolic compounds and a extraction time of 2 minutes was enough for anthocyanins.
- ❖ The developed methods presented a high repeatability and reproducibility (DER < 5%).

### REFERENCES

- Maldini M, Chessa M, Petretto GL, et al. Profiling and Simultaneous Quantitative Determination of Anthocyanins in *Wild Myrtus communis* L. Berries from Different Geographical Areas in Sardinia and their Comparative Evaluation. *Phytochem Anal.* 2016;(April):249-256.
- M. Mulas MRC. Germplasm Evaluation of Spontaneous Myrtle (*Myrtus communis* L.) for Cultivar Selection and Crop Development. *J Herbs Spices Med Plants.* 1999;6(3):31-49.
- Aleksic V, Knezevic P. Antimicrobial and antioxidative activity of extracts and essential oils of *Myrtus communis* L. *Microbiol Res.* 2014;169(4):240-254.
- Castañeda-ovando A, Pacheco-hernández MDL, Páez-hernández ME, Rodríguez JA, Galán-vidal CA. Chemical studies of anthocyanins : A review. *Food Chem.* 2009;113(4):859-871.
- Babou L, Hadidi L, Grosso C, Zaidi F, Valentão P, Andrade PB. Study of phenolic composition and antioxidant activity of myrtle leaves and fruits as a function of maturation. *European Food Research and Technology.* 2016; 242(9): 1447-1457.
- Messaoud C, Boussaid M. Myrtus communis berry color morphs: a comparative analysis o essential oils, fatty acids, phenolic compounds, and antioxidant activities. *Chem Biodiv.* 2011;8(2):300-310.